

**James Wylie, MD MHS**  
**Knee Preservation Surgeon**  
**The Orthopedic Specialty Hospital**  
**Intermountain Healthcare**  
**Office: 801-314-4900**

**KNEE ARTHROSCOPY WITH MENISCAL REPAIR**  
**PHYSICAL THERAPY PROTOCOL**

The intent of this protocol is to provide guidelines for progression of rehabilitation and is not intended to serve as a substitution for clinical decision-making. Progression through each phase of rehabilitation should take into account tissue-healing time frames, clinical objective findings, and MD approval to ensure structural stability. There will be variability between patients in terms of time frames and it is crucial not to progress through phases until the individual meets the appropriate requirements.

**INITIAL PRECAUTIONS**

**Weight Bearing:**

- 0-2 weeks→ 25% PWB
- 2-4 weeks→ 50% PWB
- 4-8 weeks→ Progress to full WBAT
- Crutch weaning and D/C is dependent walking without a limp

**Initial ROM Related Restrictions:**

- 0-90 degrees when non weight bearing
- Work on regaining terminal extension
- Hinged Knee brace locked in extension for ambulation, sleeping and activity for 2 weeks. 0-90 for exercises
- Hinged knee brace unlocked 0-90 degrees after 2 weeks. Off at night after 2 weeks.

**PHASE 1 – PROTECTION PHASE (1-4 weeks)**

**Goals:**

- Optimize bone and tissue healing and limit scar formation
- Protect surgical repair
- Reduce effusion, swelling and pain
- Restore knee ROM 0-90
- Promote normal proprioceptive and neuromuscular control

**Tissue Healing**

- PRICE – Protection, Rest, Ice, Compression, Elevation
- Scar massage after incision is healed

**Gait**

- 2 crutches at all times

POW 1-4

- Patella, patella tendon and quadriceps tendon mobilization
- Quad activation working on terminal extension
- Quad/Hamstring/Glute sets
- Straight leg raises (if no lag)
- Open chain quad exercises
- Gastrocnemius stretching without weight bearing
- Hip and Core strengthening

**PHASE 2 – PROGRESSIVE WEIGHT BEARING AND MOTION (4-8 weeks)**

**Criteria for advancement to Phase 2:**

- 0-90 ROM
- No extensor lag
- Normal joint temperature
- Minimal to no joint effusion

**Goals:**

- Progress weight bearing to full
- Normalize gait pattern on flat ground
- Progress to full symmetric ROM
- Maintain trace to no joint effusion

Precautions

- Progression of weight bearing and normal gait
- Discontinue brace at 6 weeks, can be off at night
- Closed kinetic chain activity preferred (after reaching full WB)
- No weight bearing with knee flexion greater than 90 degrees
- No cutting/pivoting/plyometrics

Strength, Proprioception and Neuromuscular Re-education

- Progressive balance training
- Calf raises
- Mini squats
- Leg presses
- Core, hip and upper body strengthening as appropriate

**PHASE 3 – EARLY STRENGTHENING (8-12 weeks)**

**Criteria for advancement to Phase 2:**

- Near symmetric ROM to contralateral side
- No extensor lag
- Normal joint temperature
- Minimal to no joint effusion

**Goals:**

- Normalize gait pattern on flat ground
- Maintain trace to no joint effusion
- Tolerate standing and walking activity

Precautions

- Progression of weight bearing and normal gait
- Closed kinetic chain activity preferred
- No cutting/pivoting/plyometrics

Strength, Proprioception and Neuromuscular Re-education

- Progressive balance training
- Calf raises
- Mini squats
- Step-ups
- Leg presses
- Double leg squats
- Core, hip and upper body strengthening as appropriate

Cardio

- Stationary Bike

### **PHASE 3 – PROGRESSIVE STRENGTHENING (12-16 weeks)**

#### **Criteria for Advancement to Phase 3:**

- Full weight bearing
- Normalize gait pattern on flat ground
- Maintain trace to no joint effusion
- Tolerate standing and walking activity

#### **Goals:**

- Build lower extremity strength, endurance, and balance.
- Quad strength 90% of contralateral side
- Normal gait/form with straight line running

#### **Precautions**

- No cutting/pivoting activities

#### **Strengthening, Proprioception and Neuromuscular Re-education**

- Double leg squats
- Leg press
- Static lunges
- Dynamic lunges
- Balance training
- Core, hip and upper body strengthening as appropriate

#### **Cardio**

- Stationary bike
- Elliptical use and treadmill walking
- Progression from treadmill walking to straight line running

## **PHASE 4 – RETURN TO SPORT (WEEKS 16+)**

### **Criteria for advancement to Phase 4**

- Good form on Phase 3 exercises
- Progressive strength/endurance development
- No joint effusion after phase 3 activities

### **Goals:**

- Build lower extremity muscular strength
- Train balance and power
- Progress to sport specific activities as tolerated

### **Precautions**

- None

### **Strengthening, Proprioception and Neuromuscular Re-education**

- Same as above:
  - Single leg squats
  - Single leg deadlifts
  - Step ups/downs
  - Multidirectional lunges
- Plyometric and cutting and pivoting exercises
  - Progression as tolerated with good form/control

### **Return to Sport Criteria**

- Full active range of motion
- No effusion
- Vertical jump, single leg hop distance, and timed single leg hop over 20 feet >85% contralateral side
- Quadriceps strength: Single leg press >90% contralateral side